


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K 31/41, 31/435, 31/46</b>		A1	(11) International Publication Number: <b>WO 98/27983</b>
			(43) International Publication Date: <b>2 July 1998 (02.07.98)</b>
(21) International Application Number: <b>PCT/DK97/00572</b> (22) International Filing Date: <b>16 December 1997 (16.12.97)</b> (30) Priority Data: 1486/96                      20 December 1996 (20.12.96)    DK (71) Applicant (for all designated States except US): <b>NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK).</b> (72) Inventors; and (75) Inventors/Applicants (for US only): <b>SAUERBERG, Per [DK/DK]; Syrenvænget 27, DK-3520 Farum (DK). HANSEN, Kristian, Tage [DK/DK]; A.D. Joergensensvej 35, 2. lejl. nr. 3, DK-2000 Frederiksberg (DK).</b> (74) Common Representative: <b>NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsvaerd (DK).</b>		(81) Designated States: <b>AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CI, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</b>  Published With international search report.	
(54) Title: <b>A METHOD OF TREATING HYPERCHOLESTEROLEMIA AND RELATED DISORDERS</b>			
(57) Abstract  The present invention relates to novel method for treating a mammal suffering from hypercholesterolemia and related disorders comprising administering to said subject an effective amount of a compound of formula (I).			
		 <div style="text-align: right;">(I)</div>	

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## A Method of Treating Hypercholesterolemia and related Disorders

### Field of the Invention

5

The present invention relates to the use of a class of compounds which are squalene synthetase inhibitors in the treatment of diseases associated with undesirable cholesterol levels in the body, and particularly diseases of the cardiovascular system, such as atherosclerosis.

10

### Background of the Invention

Hypercholesterolemia is known to be one of the prime risk factors for ischemic cardiovascular disease, such as arteriosclerosis. An important risk factor for the development of atherosclerosis is an atherogenic lipid profile i.e. hyperlipidaemia with increased LDL-cholesterol and relatively decreased HDL-cholesterol.

Typically, cholesterol is carried in the blood of warm blooded animals in certain lipid-protein complexes such as chylomicrons, very low density lipoprotein (VLDL), low density lipoprotein (LDL) and high density lipoprotein (HDL). It is widely accepted that LDL functions in a way that directly results in deposition of the LDL cholesterol in the blood vessel wall and that HDL functions in a way that results in the HDL picking up the cholesterol from the vessel wall and transporting it to the liver where it is metabolized (Brown and Goldstone, Ann. Rev. Became. 52, 223 (1983); Miller, Ann. Rev. Med. 31, 97 (1980)). It is generally accepted by those skilled in the art that reduction of abnormally high LDL (low density lipoprotein) cholesterol levels is effective therapy not only in the treatment of hyper-cholesterolemia but also in the treatment of atherosclerosis.

30, The enzyme squalene synthase (also known as squalene synthetase) is the enzyme involved in the first committed step in the de novo cholesterol biosynthesis.

---

This enzyme is a microsomal enzyme that catalyses the reductive dimerization of two molecules of farnesyl diphosphate to form squalene. Squalene is utilized only for cholesterol biosynthesis whereas farnesyl diphosphate serves as the precursor to several other biologically important compounds. The inhibition of squalene synthase  
5 would thus hinder cholesterol biosynthesis while leaving unhindered other essential pathways to e.g dolichol, ubiquinone, isopentyl tRNA and prenylated proteins.

Even though there has already been discovered squalene synthetase inhibitors a need still remains for a more effective squalene synthetase inhibitor i.e. one that provides a  
10 better antihypercholesteremic effect and exhibits a good safety profile.


We have discovered a new class of squalene synthetase inhibitors which have not previously been considered for the use in treating hypercholesterolemia and related disorders.

15

#### Summary of the Invention

The method of this invention comprises administering to a patient suffering from hypercholesterolemia and related disorders an effective amount of a compound of  
20 formula I

25



(I)

30

wherein


W is oxygen or sulphur; and

R is  $R^4$ ,  $-OR^4$ ,  $-SR^4$ ,  $-SOR^4$ ,  $-SO_2R^4$ ,  $C_{3-10}$ -cycloalkyl,  $C_{4-12}$ -(cycloalkylalkyl),  $-Z-C_{3-10}$ -cycloalkyl or  $-Z-C_{4-12}$ -(cycloalkylalkyl), wherein Z is oxygen or sulphur,  $R^4$  is  $C_{1-15}$ -alkyl,  $C_{2-15}$ -alkenyl or  $C_{2-15}$ -alkynyl, each of which is optionally substituted with one or more  
 5 halogen(s),  $-CF_3$ ,  $-CN$ , Y, phenyl or phenoxy, wherein phenyl or phenoxy is optionally substituted with halogen,  $-CN$ ,  $C_{1-4}$ -alkyl,  $C_{1-4}$ -alkoxy,  $-OCF_3$ ,  $-CF_3$ ,  $-CONH_2$  or  $-CSNH_2$ , and Y is as defined below; or

R is phenyl or benzyloxycarbonyl, each of which is optionally substituted with halogen,  $-CN$ ,  $C_{1-4}$ -alkyl,  $C_{1-4}$ -alkoxy,  $-OCF_3$ ,  $-CF_3$ ,  $-CONH_2$  or  $-CSNH_2$ ; or

- 10 R is  $-Z-Y$ ,  $-O-R^5-Z-Y$ ,  $-S-R^5-Z-Y$ ,  $-O-R^5-Z-R^4$  or  $-S-R^5-Z-R^4$  wherein Z is oxygen or sulphur,  $R^5$  is  $C_{1-15}$ -alkylene,  $C_{2-15}$ -alkenylene or  $C_{2-15}$ -alkynylene,  $R^4$  is as defined above, and Y is a 5 or 6 membered heterocyclic group containing one to four N, O or S atom(s) or a combination thereof, which heterocyclic group is optionally substituted at  
 15 carbon and/or nitrogen atom(s) with  $C_{1-6}$ -alkyl,  $-CF_3$ , phenyl, benzyl or thienyl, or a carbon atom in the heterocyclic group together with an oxygen atom form a carbonyl group, or which heterocyclic group is optionally fused with a phenyl group; and  
 G is selected from one of the following azabicyclic ring systems

20



25

- wherein  $R^1$  and  $R^2$  may be present at any position, including the point of attachment of  
 30 the  $(CH_2)_r$ -W group, and independently are hydrogen,  $C_{1-5}$ -alkyl,  $C_{2-5}$ -alkenyl,  $C_{2-5}$ -alkynyl,  $C_{1-10}$ -alkoxy,  $C_{1-5}$ -alkyl substituted with  $-OH$ , halogen,  $-NH_2$ , carboxy or phenyl; and

- R<sup>3</sup> is hydrogen, C<sub>1-6</sub>-alkyl, C<sub>2-6</sub>-alkenyl or C<sub>2-6</sub>-alkynyl; and  
n is 0, 1 or 2; and  
m is 0, 1 or 2; and  
p is 0, 1 or 2; and  
5 q is 1 or 2; and  
r is 0, 1 or 2; and  
\_\_\_\_\_ is a single or double bond; or a pharmaceutically acceptable salt thereof.

Examples of such salts include inorganic and organic acid addition salts such as  
10 hydrochloride, hydrobromide, sulphate, phosphate, acetate, fumarate, maleate, citrate,  
lactate, tartrate, oxalate, or similar pharmaceutically acceptable inorganic or organic  
acid addition salts, and include the pharmaceutically acceptable salts listed in Journal  
of Pharmaceutical Science, 66, 2 (1977) which are hereby incorporated by reference.

15 Especially preferred salts include tartrate and hydrochloride.

The terms alkyl, alkenyl, alkynyl, alkoxy, alkylene, alkenylene and alkynylene are  
intended to mean straight or branched alkyl, alkenyl, alkynyl, alkoxy, alkylene,  
alkenylene and alkynylene.

20

Typical C<sub>1-6</sub>-alkyl(ene) groups include, but are not limited to, methyl(ene), ethyl(ene), n-  
propyl(ene), iso-propyl(ene), butyl(ene), iso-butyl(ene), sec.-butyl(ene), tert.-butyl(ene),  
pentyl(ene), hexyl(ene) and the like.

25 Typical "C<sub>2-15</sub>-alkenyl(ene)" groups include but are not limited to 1-propenyl(ene), 2-  
propenyl(ene), 1,3-butadienyl(ene), 1-butenyl(ene), hexenyl(ene), pentenyl(ene) and  
the like.

Typical "C<sub>2-15</sub>-alkynyl(ene)" groups as used herein include, but are not limited to, 1-  
30 propynyl(ene), 2-propynyl(ene), 2-butylnyl(ene), 1-pentynyl(ene), 2-pentynyl(ene) and  
the like.

Typical "C<sub>1-10</sub>-alkoxy" groups include but are not limited to methoxy, ethoxy, propoxy and butoxy.

5 Typical 5 or 6 membered heterocyclic groups containing one to four N, O or S atom(s) or a combination thereof include but are not limited to pyrrole, imidazole, pyrazole, triazole, pyridine, pyrazine, pyrimidine, pyridazine, isothiazole, isoxazole, oxazole, oxadiazole, thiadiazole, quinoline, isoquinoline, quinazoline, quinoxaline, indole, benzimidazole, benzofuran, pteridine and purine.

10

As used herein, the term "halogen" means F, Cl, Br and I. Especially preferred halogens include Cl, Br and I.

15 It is to be understood that the invention extends to each of the stereoisomeric forms of the compounds of formula I as well as the racemates.

As used herein, the term patient includes any mammal which could benefit from treatment of hypercholesterolemia and related disorders. The term particularly refers to a human patient, but is not intended to be so limited.

20


The thiadiazole compounds used in the present claimed method have been disclosed and claimed in EP 0709381A1 herein incorporated by reference in its entirety. The thiadiazole compounds are known to be useful in the treatment of illnesses whose clinical manifestations are due to cholinergic deficiency.

25 Such illnesses include Alzheimer's disease, Huntingtons chorea, tardive dyskinesia, hyperkinesia, mania and Tourette Syndrome. Accordingly there is no disclosure in this reference of using the compounds to treat hypercholesterolemia and related disorders.

30 The invention also comprises the use of a compound of formula I or a pharmaceutically acceptable salt thereof for the preparation of a medicament for inhibiting squalene synthetase, especially a medicament for the treatment of hypercholesterolemia.

Preferred compounds for use in treating hypercholesterolemia and related disorders are compounds wherein G is the ring system:

5



10

and wherein m and n are 1, and p is 2.

More preferred compounds include:

15

(±)-3-Methoxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-Ethoxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-Propyloxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

20

(±)-3-Butyloxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-Pentyloxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

25

(±)-3-Hexyloxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-(4-Methylpentyloxy)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-Propylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

30

(±)-3-Butylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,



(±)-3-Pentylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(S)-3-Pentylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

5

(±)-3-Hexylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-(3,3-Dimethylbutylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

10

(±)-3-(2-(2-Thienylthio)ethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-(2,2,3,3,3-Pentafluoropropylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

15

(±)-3-(3-(2-Thienyl)propylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-Butylthio-4-((1-azabicyclo[2.2.2]octan-3-yl)methoxy)-1,2,5-thiadiazole,

20

(±)-Exo-3-pentylthio-4-(1-azabicyclo[3.2.1]octyl-6-oxy)-1,2,5-thiadiazole,

(±)-Endo-3-pentylthio-4-(1-azabicyclo[3.2.1]octyl-6-oxy)-1,2,5-thiadiazole,

(±)-Endo-3-butyloxy-4-(1-azabicyclo[2.2.1]heptyl-3-oxy)-1,2,5-thiadiazole,

25

(±)-Exo-3-butyloxy-4-(1-azabicyclo[2.2.1]heptyl-3-oxy)-1,2,5-thiadiazole,

(R)-3-Pentylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

30

(±)-3-(4-Methylpentylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

---

- (±)-3-(3-Phenylpropylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(4-Cyanobenzylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 5 (±)-3-(4-Fluorobenzylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(2-Phenylethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(2-Phenylloxyethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 10 Endo-3-butyloxy-4-(N-methyl-8-azabicyclo[3.2.1]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-Exo-3-butyloxy-4-(6-(N-methyl-8-azabicyclo[3.2.1]octan-3-onyl))-1,2,5-thiadiazole,
- 15 (±)-Endo-3-(4-cyanobenzylthio)-4-(1-azabicyclo[3.2.1]octyl-6-oxy)-1,2,5-thiadiazole,
- (±)-3-(2-(2-Thio-5-trifluoromethylthienyl)ethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 20 (±)-3-(2-(5-(2-Thienyl)thienyl)thio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(2-(2-(5-(2-Thienyl)thienyl)thio)ethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 25 (±)-3-(2-Thienylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(3-N-(2-Thiazolidonyl)propylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 30 3-Butylthio-4-(exo-2-azabicyclo[2.2.2]oct-6-yloxy)-1,2,5-thiadiazole,

Endo-3-(3-Butylthio-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,

Endo-3-(3-propylthio-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,

5 Endo-3-(3-Propylsulfonyl-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,

Endo-3-(3-(3-(4-Fluorophenyl)-2-propynyl-1-oxy)-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,

10 Endo-3-(3-(3-phenyl-2-propynyl-1-oxy)-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,

Endo-3-(3-Trifluoropropylthio-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,

15 Endo-3-(3-Trifluorobutylthio-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,

Endo-3-(3-(4-Cyanobenzylthio)-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,

Exo-3-(3-Propylthio-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[3.3.1]nonane,

20

Exo-3-(3-Butylthio-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[3.3.1]nonane,

or a pharmaceutically acceptable salt thereof.

25 Methods

a) Rat liver microsomal preparation

Male rats weighing 200 to 300 g were decapitated and the liver removed and placed in 10 ml icecold 0.25 M sucrose, 2mM EDTA and 50 mM Tris, HCl buffer pH 7.4. The

30 liver was homogenized in 3 x volume buffer relative to weight using a glass teflon Potter-Elvehjem homogenizer set to 1000 rpm. The homogenate was then centrifuged for 15 min. at 12,000 x g at 4°C. The resultant supernatant was gently decanted into 10

---

ml ultracentrifuge vials and treated at 100,000 x g for 60 min. at 4°C. The microsomal pellet was resuspended in a final volume of 1 ml icecold 0.1 M phosphate buffer pH 7.4 per gram liver tissue by gentle homogenization in a glass Potter-Elvehjem at 800 rpm. Aliquots of 0.5 ml (ca. 20 mg/ml protein) was stored at -80°C until further use in the  
5 squalene synthetase inhibition assay.

b) Assay procedure

To begin assay, 20 ml of the compound of this invention or vehicle solution is added to each 16 x 150 screw-cap culture tube on ice. Then 580 ml of N<sub>2</sub> flushed assay buffer is  
10 pipetted into each tube. 100 ml of cofactor is next added to each tube, followed by 100 ml of a dilution of microsomal enzyme (approximately 80 mg protein). The tubes are preincubated for 10 min. at 37°C, and 200 ml of the <sup>3</sup>H-FPP (200,000 dpm, 10 mM final conc.) is added to each tube at two second intervals.

15 The tubes are then incubated for exactly 10 min., shaking at 150 oscillations per min. After the 10 min. incubation, the reaction is stopped by the addition of 1 ml of 15% KOH in ethanol, and the tubes are incubated for 30 min. in a 65°C water bath for saponification of lipids and solubilization of proteins. The tubes are cooled on ice for five min. The samples are next extracted with 5 ml of petroleum ether by shaking for 10  
20 min. at low speed on a metabolic shaker. Each lower aqueous layer is frozen in a dry ice/alcohol bath (2-propanol/methanol, 1:1), and each organic layer is poured into another set of 16 x 150 screw-top culture tubes containing 2 ml of deionized water. Each ether layer is washed by vortexing each tube for 5 sec. The aqueous layers are again frozen in the dry ice/alcohol bath, and the ether is poured into scintillation vials.  
25 10 ml of AquaSol® is next added to each vial, and the vials are counted for 5 min. in a scintillation counter. Percent inhibitions are calculated from the counts obtained.

The compounds used in this method are effective over a wide dosage range. For example, in the treatment of adult humans, dosages from about 0.05 to about 100 mg,  
30 preferably from about 0.1 to about 100 mg, per day may be used. A most preferable dosage is about 10 mg to about 70 mg per day. In choosing a regimen for patients suffering from hypercholesterolemia and related disorders it may frequently be neces-

sary to begin with a dosage of from about 30 to about 70 mg per day and when the condition is under control to reduce the dosage as low as from about 1 to about 10 mg per day. The exact dosage will depend upon the mode of administration, form in which administered, the subject to be treated and the body weight of the subject to be  
5 treated, and the preference and experience of the physician or veterinarian in charge.

The route of administration may be any route, which effectively transports the active compound to the appropriate or desired site of action, such as oral or parenteral e.g. rectal, transdermal, subcutaneous, intravenous, intramuscular, topical, intranasal or an  
10 ointment, the oral route being preferred.

Typical compositions include a compound of formula I or a pharmaceutically acceptable acid addition salt thereof, associated with a pharmaceutically acceptable carrier. In making the compositions, conventional techniques for the preparation of  
15 pharmaceutical compositions may be used. For example, the active compound will usually be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a ampoule, capsule, sachet, paper, or other container. When the carrier serves as a diluent, it may be solid, semi-solid, or liquid material which acts as a vehicle, excipient, or medium for the active compound. The active  
20 compound can be adsorbed on a granular solid container for example in a sachet. Some examples of suitable carriers are water, salt solutions, alcohols, polyethylene glycols, polyhydroxyethoxylated castor oil, gelatine, lactose, amylose, magnesium stearate, talc, silicic acid, fatty acid monoglycerides and diglycerides, pentaerythritol fatty acid esters, hydroxymethylcellulose and polyvinylpyrrolidone.

25 The pharmaceutical preparations can be sterilized and mixed, if desired, with auxiliary agents, emulsifiers, salt for influencing osmotic pressure, buffers and/or coloring substances and the like, which do not deleteriously react with the active compounds.

30 For parenteral application, particularly suitable are injectable solutions or suspensions, preferably aqueous solutions with the active compound dissolved in polyhydroxylated castor oil.

---

Tablets, dragees, or capsules having talc and/or a carbohydrate carrier or binder or the like are particularly suitable for oral application. Preferable carriers for tablets, dragees, or capsules include lactose, corn starch, and/or potato starch. A syrup or elixir can be  
5 used in cases where a sweetened vehicle can be employed.

Generally, the compounds are dispensed in unit form comprising from about 1 to about 100 mg in a pharmaceutically acceptable carrier per unit dosage.

- 10 A typical tablet, appropriate for use in this method, may be prepared by conventional tableting techniques and contains:


	Active compound	5.0 mg
	Lactosum	67.8 mg Ph.Eur.
15	Avicel®	31.4 mg
	Amberlite®	1.0 mg
	Magnesii stearas	0.25 mg Ph. Eur.

CLAIMS

1. A method of inhibiting squalene synthetase in a subject in need thereof comprising administering to said subject an effective amount of a compound of formula

5 I

10



(I)

15

wherein

W is oxygen or sulphur; and

R is R<sup>4</sup>, -OR<sup>4</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, C<sub>3-10</sub>-cycloalkyl, C<sub>4-12</sub>-(cycloalkylalkyl), -Z-C<sub>3-10</sub>-cycloalkyl or -Z-C<sub>4-12</sub>-(cycloalkylalkyl), wherein Z is oxygen or sulphur, R<sup>4</sup> is C<sub>1-15</sub>-alkyl,

20 C<sub>2-15</sub>-alkenyl or C<sub>2-15</sub>-alkynyl, each of which is optionally substituted with one or more halogen(s), -CF<sub>3</sub>, -CN, Y, phenyl or phenoxy, wherein phenyl or phenoxy is optionally substituted with halogen, -CN, C<sub>1-4</sub>-alkyl, C<sub>1-4</sub>-alkoxy, -OCF<sub>3</sub>, -CF<sub>3</sub>, -CONH<sub>2</sub> or -CSNH<sub>2</sub>, and Y is as defined below; or

R is phenyl or benzyloxycarbonyl, each of which is optionally substituted with halogen,


25 -CN, C<sub>1-4</sub>-alkyl, C<sub>1-4</sub>-alkoxy, -OCF<sub>3</sub>, -CF<sub>3</sub>, -CONH<sub>2</sub> or -CSNH<sub>2</sub>; or

R is -Z-Y, -O-R<sup>5</sup>-Z-Y, -S-R<sup>5</sup>-Z-Y, -O-R<sup>5</sup>-Z-R<sup>4</sup> or -S-R<sup>5</sup>-Z-R<sup>4</sup>, wherein Z is oxygen or sulphur, R<sup>5</sup> is C<sub>1-15</sub>-alkylene, C<sub>2-15</sub>-alkenylene or C<sub>2-15</sub>-alkynylene, R<sup>4</sup> is as defined above, and Y is a 5 or 6 membered heterocyclic group containing one to four N, O or S atom(s) or a combination thereof, which heterocyclic group is optionally substituted at  
30 carbon and/or nitrogen atom(s) with C<sub>1-6</sub>-alkyl, -CF<sub>3</sub>, phenyl, benzyl or thienyl, or a carbon atom in the heterocyclic group together with an oxygen atom form a carbonyl group, or which heterocyclic group is optionally fused with a phenyl group; and

G is selected from one of the following azabicyclic ring systems

5

10



15

wherein  $R^1$  and  $R^2$  may be present at any position, including the point of attachment of the  $(CH_2)_r$ -W-group, and independently are hydrogen,  $C_{1-5}$ -alkyl,  $C_{2-5}$ -alkenyl,  $C_{2-5}$ -alkynyl,  $C_{1-10}$ -alkoxy,  $C_{1-5}$ -alkyl substituted with -OH, halogen, -NH<sub>2</sub>, carboxy or phenyl; and

20  $R^3$  is hydrogen,  $C_{1-5}$ -alkyl,  $C_{2-5}$ -alkenyl or  $C_{2-5}$ -alkynyl; and

$n$  is 0, 1 or 2; and

$m$  is 0, 1 or 2; and

$p$  is 0, 1 or 2; and

$q$  is 1 or 2; and

25  $r$  is 0, 1 or 2; and

..... is a single or double bond;


or a pharmaceutically acceptable salt thereof.

2. A method of inhibiting squalene synthetase in a subject in need thereof  
30 comprising administering to said subject an effective amount of a compound of formula I

I



5



(I)

10

wherein

W is oxygen or sulphur; and

R is  $R^4$ ,  $-OR^4$ ,  $-SR^4$ ,  $-SOR^4$ ,  $-SO_2R^4$ ,  $C_{3-10}$ -cycloalkyl,  $C_{4-12}$ -(cycloalkylalkyl),  $-Z-C_{3-10}$ -cycloalkyl or  $-Z-C_{4-12}$ -(cycloalkylalkyl), wherein Z is oxygen or sulphur,  $R^4$  is  $C_{1-15}$ -alkyl,

15  $C_{2-15}$ -alkenyl or  $C_{2-15}$ -alkynyl, each of which is substituted with one or more halogen(s),  $-CF_3$ ,  $-CN$ , Y, phenyl or phenoxy, wherein phenyl or phenoxy is optionally substituted with halogen,  $-CN$ ,  $C_{1-4}$ -alkyl,  $C_{1-4}$ -alkoxy,  $-OCF_3$ ,  $-CF_3$ ,  $-CONH_2$  or  $-CSNH_6$ , and Y is as defined below; or

R is phenyl or benzyloxycarbonyl, each of which is optionally substituted with halogen,

20  $-CN$ ,  $C_{1-4}$ -alkyl,  $C_{1-4}$ -alkoxy,  $-OCF_3$ ,  $-CF_3$ ,  $-CONH_2$  or  $-CSNH_2$ ; or

R is  $-Z-Y$ ,  $-O-R^5-Z-Y$ ,  $-S-R^5-Z-Y$ ,  $-O-R^5-Z-R^4$  or  $-S-R^5-Z-R^4$ , wherein Z is oxygen or sulphur,  $R^5$  is  $C_{1-15}$ -alkylene,  $C_{2-15}$ -alkenylene or  $C_{2-15}$ -alkynylene,  $R^4$  is as defined





above, and Y is a 5 or 6 membered heterocyclic group containing one to four N, O or S atom(s) or a combination thereof, which heterocyclic group is optionally substituted at

25 carbon and/or nitrogen atom(s) with  $C_{1-6}$ -alkyl,  $-CF_3$ , phenyl, benzyl or thienyl, or a carbon atom in the heterocyclic group together with an oxygen atom form a carbonyl group, or which heterocyclic group is optionally fused with a phenyl group; and

G is selected from one of the following azabicyclic ring systems

30

5



10

wherein  $R^1$  and  $R^2$  may be present at any position, including the point of attachment of the  $(CH_2)_r$ -W-group, and independently are hydrogen,  $C_{1-5}$ -alkyl,  $C_{2-5}$ -alkenyl,  $C_{2-5}$ -alkynyl,  $C_{1-10}$ -alkoxy,  $C_{1-5}$ -alkyl substituted with -OH, halogen,  $-NH_2$ , carboxy or phenyl;

15

and

$R^3$  is hydrogen,  $C_{1-5}$ -alkyl,  $C_{2-5}$ -alkenyl or  $C_{2-5}$ -alkynyl; and

$n$  is 0, 1 or 2; and

$m$  is 0, 1 or 2; and

$p$  is 0, 1 or 2; and

20

$q$  is 1 or 2; and

$r$  is 0, 1 or 2; and


..... is a single or double bond;

or a pharmaceutically acceptable salt thereof.

25

3. The method according to claim 1 or 2 wherein G in formula I is the ring system

30



wherein R<sup>1</sup> and R<sup>2</sup> are as defined in claim 1, and m and n are 1, and p is 2;  
or a pharmaceutically acceptable salt thereof.

- 5     4.     The method according to anyone of the preceding claims wherein the  
compound is selected from the following:

(±)-3-Methoxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

10     (±)-3-Ethoxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-Propyloxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-Butyloxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

15

(±)-3-Pentyloxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-Hexyloxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

20     (±)-3-(4-Methylpentyloxy)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-Propylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-Butylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

25

(±)-3-Pentylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(S)-3-Pentylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

30

(±)-3-Hexylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

---

- (±)-3-(3,3-Dimethylbutylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 5 (±)-3-(2-(5-(2-Thienyl)thienyl)thio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(2-(2-Thienylthio)ethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(2-Thienylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 10 (±)-3-(2,2,3,3,3-Pentafluoropropylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(3-(2-Thienyl)propylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 15 (±)-3-Butylthio-4-((1-azabicyclo[2.2.2]octan-3-yl)methoxy)-1,2,5-thiadiazole,
- (±)-Exo-3-pentylthio-4-(1-azabicyclo[3.2.1]octyl-6-oxy)-1,2,5-thiadiazole,
- 20 (±)-Endo-3-pentylthio-4-(1-azabicyclo[3.2.1]octyl-6-oxy)-1,2,5-thiadiazole,
- (±)-Endo-3-butylthio-4-(1-azabicyclo[2.2.1]heptyl-3-oxy)-1,2,5-thiadiazole,
- (±)-Exo-3-butylthio-4-(1-azabicyclo[2.2.1]heptyl-3-oxy)-1,2,5-thiadiazole,
- 25 (R)-3-Pentylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(4-Methylpentylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 30 (±)-3-(3-Phenylpropylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

- (±)-3-(4-Cyanobenzylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(4-Fluorobenzylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 5 (±)-3-(2-Phenylethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(2-Phenyloxyethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- Endo-3-butyloxy-4-(N-methyl-8-azabicyclo[3.2.1]octyl-3-oxy)-1,2,5-thiadiazole,
- 10 (±)-Exo-3-butyloxy-4-(6-(N-methyl-8-azabicyclo[3.2.1]octan-3-onyl))-1,2,5-thiadiazole,
- (±)-Endo-3-(4-cyanobenzylthio)-4-(1-azabicyclo[3.2.1]octyl-6-oxy)-1,2,5-thiadiazole,
- 15 (±)-3-(2-(2-Thio-5-trifluoromethylthienyl)ethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(2-(2-(5-(2-Thienyl)thienyl)thio)ethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 20 (±)-3-(3-N-(2-Thiazolidonyl)propylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 3-Butylthio-4-(exo-2-azabicyclo[2.2.2]oct-6-yloxy)-1,2,5-thiadiazole,
- 25 Endo-3-(3-Butylthio-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,
- Endo-3-(3-propylthio-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,
- 30 Endo-3-(3-Propylsulfonyl-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,

Endo-3-(3-(3-(4-Fluorophenyl)-2-propynyl-1-oxy)-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,

5 Endo-3-(3-(3-phenyl-2-propynyl-1-oxy)-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,

Endo-3-(3-Trifluoropropylthio-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,

10 Endo-3-(3-Trifluorobutylthio-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,

Endo-3-(3-(4-Cyanobenzylthio)-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[2.2.1]heptane,

Exo-3-(3-Propylthio-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[3.3.1]nonane,

15 Exo-3-(3-Butylthio-1,2,5-thiadiazol-4-yloxy)-1-azabicyclo[3.3.1]nonane,

or a pharmaceutically acceptable salt thereof.

5. The method according to anyone of the preceding claims wherein the  
20 compound is selected from the following:

(±)-3-Methoxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-Ethoxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

25

(±)-3-Propyloxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-Butyloxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

30 (±)-3-Pentyloxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

- (±)-3-Hexyloxy-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(4-Methylpentyloxy)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 5 (±)-3-Propylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-Butylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-Pentylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 10 (S)-3-Pentylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-Hexylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 15 (±)-3-(3,3-Dimethylbutylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(2-(5-(2-Thienyl)thienyl)thio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(2-(2-Thienylthio)ethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 20 (±)-3-(2-Thienylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-(2,2,3,3,3-Pentafluoropropylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- 25 (±)-3-(3-(2-Thienyl)propylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,
- (±)-3-Butylthio-4-((1-azabicyclo[2.2.2]octan-3-yl)methoxy)-1,2,5-thiadiazole,
- 30 (R)-3-Pentylthio-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-(4-Methylpentylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-(3-Phenylpropylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

5 (±)-3-(4-Cyanobenzylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-(4-Fluorobenzylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-(2-Phenylethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

10

(±)-3-(2-Phenyloxyethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-(2-(2-Thio-5-trifluoromethylthienyl)ethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

15

(±)-3-(2-(2-(5-(2-Thienyl)thienyl)thio)ethylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-1,2,5-thiadiazole,

(±)-3-(3-N-(2-Thiazolidonyl)propylthio)-4-(1-azabicyclo[2.2.2]octyl-3-oxy)-

20

1,2,5-thiadiazole,

3-Butylthio-4-(exo-2-azabicyclo[2.2.2]oct-6-yloxy)-1,2,5-thiadiazole,

or a pharmaceutically acceptable salt thereof.

25

6. A method of treating hypercholesterolemia in a subject in need thereof comprising administering to said subject an effective amount of a compound according to claim 1.

30

7. A method of treating hypercholesterolemia in a subject in need thereof comprising administering to said subject an effective amount of a compound according to claim 2.



8. A method of treating hypercholesterolemia in a subject in need thereof comprising administering to said subject an effective amount of a compound according to claim 3.

5

9. A method of treating hypercholesterolemia in a subject in need thereof comprising administering to said subject an effective amount of a compound according to claim 4.

10 10. A method of treating hypercholesterolemia in a subject in need thereof comprising administering to said subject an effective amount of a compound according to claim 5.

11. The use of a compound according to claim 1 or a pharmaceutically acceptable salt thereof for the preparation of a medicament for inhibiting squalene synthetase.

12. The use of a compound according to claim 1 or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treatment of hypercholesterolemia.

20 13. The use of a compound according to claim 2 or a pharmaceutically acceptable salt thereof for the preparation of a medicament for inhibiting squalene synthetase.

14. The use of a compound according to claim 2 or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treatment of hypercholesterolemia.

25

15. The use of a compound according to claim 3 or a pharmaceutically acceptable salt thereof for the preparation of a medicament for inhibiting squalene synthetase.

16. The use of a compound according to claim 3 or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treatment of hypercholesterolemia.

30

17. The use of a compound according to claim 4 or a pharmaceutically acceptable salt thereof for the preparation of a medicament for inhibiting squalene synthetase.

18. The use of a compound according to claim 4 or a pharmaceutically acceptable  
5 salt thereof for the preparation of a medicament for  
treatment of hypercholesterolemia.

19. The use of a compound according to claim 5 or a pharmaceutically acceptable  
10 salt thereof for the preparation of a medicament for inhibiting squalene synthetase.

20. The use of a compound according to claim 5 or a pharmaceutically acceptable  
15 salt thereof for the preparation of a medicament for  
treatment of hypercholesterolemia.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**